

Hamilton-Jacobi formulation of systems within Caputo's fractional derivative

February 2, 2008

Eqab M.Rabei¹

Department of Physics, Mutah University, Karak, Jordan

Ibtesam Almayteh

Department of Physics, Mutah University, Karak, Jordan

Sami I. Muslih²

Department of Physics, Al-Azhar University, Gaza, Palestine

Dumitru Baleanu³

Department of Mathematics and Computer Sciences, Faculty of Arts and
Sciences, Çankaya University- 06530, Ankara, Turkey

Keywords: fractional derivative, Hamilton-Jacobi formulation

PACS:

Abstract

In this paper we develop a fractional Hamilton-Jacobi formulation for discrete systems in terms of fractional Caputo derivatives. The fractional action function is obtained and the solutions of the equations of motion are recovered. An example is studied in details.

¹E-mail: eqabrabei@yahoo.com

²E-mail: smuslih@ictp.trieste.it

³On leave of absence from Institute of Space Sciences, P.O.BOX, MG-23, R 76900, Magurele-Bucharest, Romania, E-mails: dumitru@cankaya.edu.tr, baleanu@venus.nipne.ro

1 Introduction

Fractional calculus generalized the classical calculus and it has many important applications in various fields of science and engineering [1-10]. These applications include classical and quantum mechanics, field theory, and optimal control [11-30] formulated mostly in terms of Riemann-Liouville (RL) and Caputo fractional derivatives. In contrast with RL derivative, Caputo derivative of a constant is zero, and for a fractional differential equation defined in terms of Caputo derivatives the standard boundary conditions are well defined. Therefore, this kind of fractional derivative gained importance among engineers and scientists.

Recently, the fractional Hamiltonian formulations were investigated for fractional discrete and continuous systems in terms of RL and Caputo derivatives [22-37]. Even more recently, a fractional Hamilton-Jacobi formalism within RL derivative was proposed in [38].

As it well known the classical HamiltonJacobi equation represents a reformulation of classical mechanics which equivalent to other formulations such as Newton's laws of motion, Lagrangian mechanics and Hamiltonian mechanics. In addition, the HamiltonJacobi equation is useful in finding the conserved quantities for mechanical systems, which may be possible even when the mechanical problem itself cannot be solved completely. The Hamilton-Jacobi theory represents the only formulation of mechanics in which the motion of a particle can be represented as a wave.

On the other hand in the area of fractional mechanics the investigation of the fractional Hamilton-Jacobi equation is still at the beginning of its development.

Due to the above mentioned reasons the formulation of a fractional Hamilton-Jacobi within Caputo fractional derivatives is an interesting issue to be investigated.

The plan of this paper is as follows:

In section two the basic mathematical tools are briefly described. In section three the fractional Hamilton equations within Caputo's derivative are mentioned. Section four contains the fractional Hamilton-Jacobi formulation by using Caputo's derivative. An illustrative example is analyzed in section five. Finally, section six contains our conclusions.

2 Basic definitions

In the following we briefly present some fundamental definitions used in the previous section.

The left and the right Riemann-Liouville and Caputo fractional derivatives are defined as follows:

The Left Riemann-Liouville Fractional Derivative

$${}_a D_t^\alpha f(t) = \frac{1}{\Gamma(n-\alpha)} \left(\frac{d}{dt} \right)^n \int_a^t (t-\tau)^{n-\alpha-1} f(\tau) d\tau, \quad (1)$$

The Right Riemann-Liouville Fractional Derivative

$${}_t D_b^\alpha f(t) = \frac{1}{\Gamma(n-\alpha)} \left(-\frac{d}{dt} \right)^n \int_t^b (\tau-t)^{n-\alpha-1} f(\tau) d\tau, \quad (2)$$

The corresponding Caputo's fractional derivatives are defined as follows:

The Left Caputo Fractional Derivative

$${}_a^C D_t^\alpha f(t) = \frac{1}{\Gamma(n-\alpha)} \int_a^t (t-\tau)^{n-\alpha-1} \left(\frac{d}{d\tau} \right)^n f(\tau) d\tau, \quad (3)$$

and

The Right Caputo Fractional Derivative

$${}_t^C D_b^\alpha f(t) = \frac{1}{\Gamma(n-\alpha)} \int_t^b (\tau-t)^{n-\alpha-1} \left(-\frac{d}{d\tau} \right)^n f(\tau) d\tau, \quad (4)$$

where α represents the order of the derivative such that $n-1 < \alpha < n$.

In [22] the fractional Euler-Lagrange equations are obtained. We present briefly the main result obtained in the following (for more details see [22]).

Let $J[q]$ be a functional of the form

$$J[q] = \int_a^b L(t, q, {}_a^C D_t^\alpha q, {}_t^C D_b^\beta q) dt \quad (5)$$

where $0 < \alpha, \beta < 1$ and defined on the set of functions $y(x)$ which have continuous LCFD of order α and RCFD of order β in $[a, b]$. Then a necessary

condition for $J[q]$ to have an extremum for a given function $q(t)$ is that $q(t)$ satisfy the generalized Euler-Lagrange equation given by

$$\frac{\partial L}{\partial q} + {}_tD_b^\alpha \frac{\partial L}{\partial {}_a^C D_t^\alpha q} + {}_aD_t^\beta \frac{\partial L}{\partial {}_t^C D_b^\beta q} = 0, \quad t \in [a, b] \quad (6)$$

and the transversality conditions given by

$$\left[{}_tD_b^{\alpha-1} \left(\frac{\partial L}{\partial {}_a^C D_t^\alpha q} \right) - {}_aD_t^{\beta-1} \left(\frac{\partial L}{\partial {}_t^C D_b^\beta q} \right) \right] \eta(t)|_a^b = 0. \quad (7)$$

3 Fractional Hamiltonian formulation

In this section we briefly present the Hamiltonian formulation within Caputo's fractional derivatives.

Let us consider the fractional Lagrangian as given below

$$L(q, {}_a^C D_t^\alpha q, {}_t^C D_b^\beta q, t), \quad 0 < \alpha, \beta < 1. \quad (8)$$

By using (8) we define the canonical momenta p_α and p_β as follows

$$p_\alpha = \frac{\partial L}{\partial {}_a^C D_t^\alpha q}, \quad p_\beta = \frac{\partial L}{\partial {}_t^C D_b^\beta q}. \quad (9)$$

Making use of (8) and (9) we define the fractional canonical Hamiltonian as

$$H = p_\alpha {}_a^C D_t^\alpha q + p_\beta {}_t^C D_b^\beta q - L. \quad (10)$$

The fractional Hamilton equations are obtained as follows [37]

$$\frac{\partial H}{\partial t} = -\frac{\partial L}{\partial t}, \quad \frac{\partial H}{\partial p_\alpha} = {}_a^C D_t^\alpha q, \quad \frac{\partial H}{\partial p_\beta} = {}_t^C D_b^\beta q, \quad \frac{\partial H}{\partial q} = {}_tD_b^\alpha p_\alpha + {}_aD_t^\beta p_\beta. \quad (11)$$

4 Fractional Hamilton-Jacobi formulation

In this section, we construct the Hamilton-Jacobi formulation within Caputo fractional derivatives. According to [37], the fractional Hamiltonian is defined in equation (10). Now, let consider the canonical transformations

with the generating function $F_2({}_a^C D_t^{\alpha-1} q, {}_t^C D_b^{\beta-1} q, P_\alpha, P_\beta, t) = S$. The new Hamiltonian will take the form

$$K = P_\alpha {}_a^C D_t^\alpha Q + P_\beta {}_t^C D_b^\beta Q - L(Q, {}_a^C D_t^\alpha Q, {}_t^C D_b^\beta Q). \quad (12)$$

The variations of the following integrals

$$\delta \int_{t_1}^{t_2} (P_\alpha {}_a^C D_t^\alpha Q + P_\beta {}_t^C D_b^\beta Q - K) dt = 0, \quad \delta \int_{t_1}^{t_2} (p_\alpha {}_a^C D_t^\alpha q + p_\beta {}_t^C D_b^\beta q - H) dt = 0, \quad (13)$$

are vanishing, then we obtain the relation between the Hamiltonians H and K as follows

$$p_\alpha {}_a^C D_t^\alpha q + p_\beta {}_t^C D_b^\beta q - H = P_\alpha {}_a^C D_t^\alpha Q + P_\beta {}_t^C D_b^\beta Q - K + \frac{dF}{dt}, \quad (14)$$

where the function F is given as

$$F = S({}_a^C D_t^{\alpha-1} q, {}_t^C D_b^{\beta-1} q, P_\alpha, P_\beta, t) - P_\alpha {}_a^C D_t^{\alpha-1} Q - P_\beta {}_t^C D_b^{\beta-1} Q. \quad (15)$$

Substituting this function F in equation (14), we obtain the following equations

$${}_a^C D_t^{\alpha-1} Q = \frac{\partial S}{\partial P_\alpha}, \quad {}_t^C D_b^{\beta-1} Q = \frac{\partial S}{\partial P_\beta} \quad (16)$$

$$p_\alpha = \frac{\partial S}{\partial {}_a^C D_t^{\alpha-1} q}, \quad p_\beta = \frac{\partial S}{\partial {}_t^C D_b^{\beta-1} q}, \quad (17)$$

$$K = H + \frac{\partial S}{\partial t}. \quad (18)$$

In the case that the new variables (Q, P_α, P_β) are constants in time, then $K = 0$. One can easily show that action function can be put in the form.

$$S = \int_{t_1}^{t_2} L dt. \quad (19)$$

For time-independent Hamiltonian H , the action function S can be put in the form

$$S = W_1({}_a^C D_t^{\alpha-1} q, E_1) + W_2({}_t^C D_b^{\beta-1} q, E_2) + f(E_1, E_2, t), \quad (20)$$

here, W is called the Hamilton's characteristic function and $f(E_1, E_2, t) = -Et$, where $E_1 = P_\alpha$, $E_2 = P_\beta$ and $E = E_1 + E_2$.

Making use of equation (22), and after some simple manipulations, we obtain

$${}_a^C D_t^{\alpha-1} Q = \frac{\partial S}{\partial P_\alpha} = \lambda_1, \quad {}_t^C D_b^{\beta-1} Q = \frac{\partial S}{\partial P_\beta} = \lambda_2 \quad (21)$$

$$p_\alpha = \frac{\partial W_1}{\partial {}_a^C D_t^{\alpha-1} q}, \quad p_\beta = \frac{\partial W_2}{\partial {}_t^C D_b^{\beta-1} q}, \quad (22)$$

$$\frac{\partial S}{\partial t} = -H = -E. \quad (23)$$

and the Hamilton-Jacobi partial differential equation for fractional system reads as

$$\frac{\partial S}{\partial t} + H = 0. \quad (24)$$

Now the solutions of equation (20), give the values of W_1 and W_2 as

$$W_1 = \int p_\alpha d {}_a^C D_t^{\alpha-1} q, \quad W_2 = \int p_\beta d {}_t^C D_b^{\beta-1} q. \quad (25)$$

5 Illustrative example

In this section, we shall present an example to demonstrate the application of the formulation developed in the previous section. Let us consider the following Hamiltonian [36]

$$H = \frac{1}{2} p_\alpha^2 + q. \quad (26)$$

The fractional Hamilton-Jacobi equation is given by

$$\frac{1}{2} p_\alpha^2 + q - E_1 = 0. \quad p_\beta = 0. \quad (27)$$

Making use of equation (22), we get

$$\left[\frac{\partial W_1}{\partial {}_a^C D_t^{\alpha-1} q} \right]^2 + q - E_1 = 0, \quad (28)$$

which has the following solution

$$W_1 = \sqrt{2(E_1 - q)} {}_a^C D_t^{\alpha-1} q, \quad (29)$$

this equation leads us to obtain p_α and S respectively as

$$p_\alpha = \sqrt{2(E_1 - q)} \quad (30)$$

$$S = \sqrt{2(E_1 - q)} {}^C D_t^{\alpha-1} q - E_1 t. \quad (31)$$

Again making use of equations (21), (31), we arrive at

$${}_a^C D_t^{\alpha-1} Q = \frac{\partial S}{\partial t} = \frac{1}{\sqrt{2(E_1 - q)}} {}^C D_t^{\alpha-1} q - t = \lambda_1. \quad (32)$$

Equation (32), can be easily solved to obtain

$${}_a^C D_t^{\alpha-1} q = \sqrt{2(E_1 - q)}(t + \lambda_1), \quad (33)$$

or

$${}_a^C D_t^\alpha q = \sqrt{2(E_1 - q)} = p_\alpha. \quad (34)$$

Taking the Riemman-Liouville derivative of equation (34), we obtain

$${}_t D_b^{\beta C} {}_a^C D_t^\alpha q = 1. \quad (35)$$

This result is in exact agreement with that obtained if we use the fractional Hamiltonian formulation.

6 Conclusions

In this paper, we have obtained the Hamilton-Jacobi partial differential equation within Caputo's fractional derivative. Finding the action function leads us to obtain the solutions of equations of motion. An example was investigated and the solutions of the equations of motion are in exact agreement with those obtained by using the Hamiltonian formulation. The advantages of using the method presented in this paper, is that we can easily obtain the action function, which is the essential part to obtain the path integral quantization for any mechanical fractional system, and this topic is now under investigation.

References

- [1] Oldham K B and Spanier J 1974 *The Fractional Calculus*, Academic Press, New-York.
- [2] Miller K S and Ross B 1993 *An Introduction to the Fractional Integrals and Derivatives-Theory and Applications*, John Wiley and Sons Inc., New York.
- [3] Samko S G, Kilbas A A and Marichev O I 1993 *Fractional Integrals and Derivatives - Theory and Applications*, Gordon and Breach, Linghorne, P.A.
- [4] Hilfer R 2000 *Applications of Fractional Calculus in Physics*, World Scientific Publishing Company, Singapore.
- [5] Podlubny I 1999 *Fractional Differential Equations*, Academic Press, San Diego CA.
- [6] Zaslavsky G M 2005 *Hamiltonian Chaos and Fractional Dynamics*, Oxford University Press, Oxford.
- [7] Kilbas, A A, Srivastava H M and Trujillo J J 2006 *Theory and Applications of Fractional Differential Equations*, Elsevier.
- [8] Magin R L 2006 *Fractional Calculus in Bioengineering*, Begell House Publisher, Inc. Connecticut.
- [9] Mainardi F, Luchko Y and Pagnini G 2001 *Frac. Calc. Appl. Anal.* **4(2)** 153.
- [10] Naber M 2004 *J. Math. Phys.* **45** 3339.
- [11] Riewe F 1996 *Phys. Rev. E* **53** 1890.
- [12] Riewe F 1997 *Phys. Rev. E* **55** 3581.
- [13] Klimek M 2001 *Czech. J. Phys.* **51** 1348.
- [14] Klimek M 2002 *Czech. J. Phys.* **52** 1247.
- [15] Klimek M 2005 *Czech. J. Phys.* **55** 1447.

- [16] Agrawal O P 2002 *J. Math. Anal. Appl.* **272** 368.
- [17] Dreisigmeyer D W and Young P M 2003 *J. of Phys. A: Math. Gen.* **36** 8297.
- [18] Dreisigmeyer D W and Young P M 2004 *J. of Phys. A: Math. Gen.* **37** L117.
- [19] Tofighi A 2003 *Physica A* **329** 29.
- [20] Tofighi A and Pour H N 2007 *Physica A* **374(1)** 41.
- [21] Baleanu D 2004 *Constrained systems and Riemann-Liouville fractional derivative*, Proceedings of 1st IFAC Workshop on Fractional Differentiation and its Applications(Bordeaux, France, July 19-21, p. 597).
- [22] Agrawal O P 2007 Generalized Euler-Lagrange equations and transversality conditions for FVPs in terms of Caputo Derivative, in *Proc. MME06, Ankara, Turkey, April 27-29, 2006* (Eds. K. Tas, J.A. Tenreiro Machado and D. Baleanu), to appear in *J. Vib. Contr.*
- [23] Rabei E M, Nawafleh K I, Hijjawi R S, Muslih S I and Baleanu D, 2007 *J. Math. Anal. Appl.* **327** 891.
- [24] Momani S , Qaralleh R 2006 *Comp.Math.Appl.* **52** 459.
- [25] Baleanu D and Muslih S 2005 New trends in fractional quantization method. *Intelligent Systems at the Service of Mankind*, volume 2. U-Books Verlag, Augsburg, Germany, 1st edition.
- [26] Baleanu D 2005 Constrained systems and Riemann-Liouville fractional derivatives. *Fractional differentiation and its applications*. U-Books Verlag, Augsburg, Germany.
- [27] Muslih S I and Baleanu D 2005 *J. Math. Anal. Appl.* **304** 599.
- [28] Baleanu D 2006 *Signal Processing* **86(10)** 2632.
- [29] Muslih S I and Baleanu D 2005 *Czech. J. Phys.* **55(6)** 633.
- [30] Baleanu D and Muslih S I 2005 *Physica Scripta* **72(2-3)** 119.

- [31] Agrawal O P 2006 A Formulation and a Numerical Scheme for Fractional Optimal Control Problems, *in Proceedings of the IFAC/FDA06, Porto, Portugal, July 19-21.*
- [32] Agrawal O P 2004 Nonlinear Dynamics **38** 323.
- [33] Agrawal O P and Baleanu D 2006 A Hamiltonian Formulation and a Direct Numerical Scheme for Fractional Optimal Control Problems, *in Proc. MME06, Ankara, Turkey, April 27-29* (Eds. K. Tas, J.A. Tenreiro Machado and D. Baleanu), to appear in *J. Vib. Contr.* (2007).
- [34] Rabei E M Ajlouni A W and Ghassib, H B 2006 *Int. J. Theor. Phys.*, 45,1619. Rabei E M Ajlouni A W and Ghassib, H B 2006 WSEA Transaction on Mathematics **5** 853.
- [35] Rabei E M, Al-Halholi T and Rousan A 2004 *Int. J. Mod. Phy. A* **19** 3083.
- [36] Agrawal O P 2007 Generalized Euler- lagrange equations and transversality conditions for FVPs in terms of Caputo Derivative, in Proc. MME06, Ankara, Turkey, April 27-29, 2006, to appear in *J. Vib. Contr.*
- [37] Baleanu D and Agrawal O P 2006 *Czech. J. Phys.*, 56(10-11) 1087, arXiv;math-ph/0612025 v1 7 Dec 2006.
- [38] Rabei E M and Ababenh B S 2007 *J. Math. Anal. Appl.*, (in press)